

§4.4b The Second Fundamental Theorem of Calculus
The Second Fundamental Theorem of Calculus

Notes based on: *Calculus for AP* by Larson & Battaglia. © 2017 Cengage Learning.
Calculus, AP Edition, 9th ed. by Larson & Edwards. © 2010 Brooks/Cole, Cengage Learning.

Learning Goals: Students will be able to...

- Understand and use the Second Fundamental Theorem of Calculus.

Learning Objectives: Students will be able to...


- 2.1C Calculate derivatives.
- 2.1D Determine higher order derivatives.
- 2.2A Use derivatives to analyze properties of a function.
- 3.2C Calculate a definite integral using areas and properties of definite integrals.
- 3.3A Analyze functions defined by an integral.

The Second Fundamental Theorem of Calculus


Earlier, we saw that the definite integral of f on the interval $[a, b]$ as defined using the constant b as the upper limit of integration and x as the variable of integration.

However, a slightly different situation may arise in which the variable x is used in the upper limit of integration. To avoid the confusion of using x in two different ways, t is temporarily used as the variable of integration.

The Definite Integral as a Number



The Definite Integral as a Function of x



The Second Fundamental Theorem of Calculus

Suppose $F(x) = \int_0^x \cos(t) dt$.

If we attempt to evaluate this definite integral using the Fundamental Theorem of Calculus:

$$\begin{aligned} F(x) &= \int_0^x \cos(t) dt = [\sin(t)]_{t=0}^{t=x} \\ &= \sin(x) - \sin(0) = \sin(x) - 0 \\ &= \sin(x) \end{aligned}$$

Note that the derivative of F is the original integrand (with only the variable changed). That is:

$$\frac{d}{dx}[F(x)] = \frac{d}{dx}[\sin(x)] = \frac{d}{dx}\left[\int_0^x \cos(t) dt\right] = \cos(x)$$

This result is generated in the next theorem, called the **Second Fundamental Theorem of Calculus**.

The Second Fundamental Theorem of Calculus

THEOREM THE SECOND FUNDAMENTAL THEOREM OF CALCULUS

If f is continuous on an open interval I containing a , then, for every x in the interval,

$$\frac{d}{dx}\left[\int_a^x f(t) dt\right] = f(x).$$

Two things are necessary to utilize this theorem:

- (1) The lower limit of integration must be a constant value.
- (2) The upper limit of integration must be x .

The Second Fundamental Theorem of Calculus

Otherwise, we must use a more generalized version of this theorem.

Suppose $F(x) = \int_a^{g(x)} f(t) dt$. If we say that $u = g(x)$, then $F(x) = \int_a^u f(t) dt$.

Utilizing the Chain Rule, we get $F'(x) = f(u) u' = f(g(x)) \cdot g'(x)$.

Thus, a more generalized version of the Second Fundamental Theorem of Calculus states:

$$\frac{d}{dx}\left[\int_a^{g(x)} f(t) dt\right] = f(g(x)) \cdot g'(x)$$

Example: The Second Fundamental Theorem of Calculus


Find $F'(x)$ if $F(x) = \int_{\pi^2}^{x^6} \sqrt[3]{t} dt$.

Example: The Second Fundamental Theorem of Calculus

The graph of the function f shown consists of a semicircle and three line segments. Let g be the function given by

$$g(x) = \int_{-2}^x f(t) dt.$$

(a) Find $g(3)$, $g'(3)$, and $g''(3)$.




Example: The Second Fundamental Theorem of Calculus

The graph of the function f shown consists of a semicircle and three line segments. Let g be the function given by

$$g(x) = \int_{-2}^x f(t) dt.$$

(b) Find all values of x in the open interval $(-4, 6)$ at which the graph of g is increasing and concave downward. Justify your answer.




Example: The Second Fundamental Theorem of Calculus

The graph of the function f shown consists of a semicircle and three line segments. Let g be the function given by

$$g(x) = \int_{-2}^x f(t) dt.$$

(c) Find all values of x in the open interval $(-4, 6)$ at which g attains a relative minimum. Justify your answer.




Example: The Second Fundamental Theorem of Calculus

The graph of the function f shown consists of a semicircle and three line segments. Let g be the function given by

$$g(x) = \int_{-2}^x f(t) dt.$$

(d) Find the absolute minimum value of g on the closed interval $[-4, 6]$. Justify your answer.



Example: The Second Fundamental Theorem of Calculus

The graph of the function f shown consists of a semicircle and three line segments. Let g be the function given by

$$g(x) = \int_{-2}^x f(t) dt.$$

(e) Find all values of x in the open interval $(-4, 6)$ at which the graph of g has a point of inflection. Justify your answer.

